

Brief Clinical Report

Nasal Dimple as Part of the 22q11.2 Deletion Syndrome

Karen W. Gripp,^{1*} Donna M. McDonald-McGinn,^{1,4} Deborah A. Driscoll,^{1,2,3} Lori A. Reed,¹ Beverly S. Emanuel,^{1,3} and Elaine H. Zackai^{1,3,4}

¹Division of Human Genetics and Molecular Biology, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania

²Division of Reproductive Genetics, University of Pennsylvania School of Medicine, Philadelphia

³Department of Pediatrics, University of Pennsylvania School of Medicine, Philadelphia

⁴Department of Obstetrics and Gynecology, University of Pennsylvania School of Medicine, Philadelphia

The phenotype of the 22q11.2 microdeletion syndrome is quite variable. We describe 2 patients with a 22q11.2 deletion and a dimpled nasal tip, which, we suggest can be the extreme of the broad or bulbous nose commonly found in the 22q11.2 deletion syndrome, and should not be confused with the more severe nasal abnormalities seen in frontonasal dysplasia. Am. J. Med. Genet. 69:290–292, 1997. © 1997 Wiley-Liss, Inc.

KEY WORDS: DiGeorge syndrome; conotruncal anomaly face syndrome (CAFS); nasal dimple; velo-cardio-facial syndrome (VCFS); 22q11.2 deletion

PATIENT 1

A caucasian female was born vaginally at term to a 23-year-old primigravida and a 25-year-old non-consanguineous father. The pregnancy was complicated by light bleeding in the first trimester and elevated maternal blood pressure at 33 weeks of gestation. Birth weight was 2.76 kg (10–25th centile); length 47 cm (10–25th centile); OFC 33 cm (25–50th centile). At 8 days the infant was found to have a broad nose with some lateral build-up and a dimple in the nasal tip (Fig. 1), a nevus flammeus over the left upper eyelid, an inner canthal distance of 2.5 cm, outer canthal distance of 6 cm, and interpupillary distance of 4 cm (50th centile). Ears appeared “squared off” with overfolded helices. The mouth was well formed and the palate grossly in-

tact. A 2/6 systolic heart murmur and an umbilical hernia were present. Female external genitalia had a cleft appearance of the anterior labia majora. The limbs and dermatoglyphics were normal.

Echocardiography showed a tetralogy of Fallot (TOF). Chromosomal studies demonstrated a 46,XX karyotype. Because of the association of 22q11.2 microdeletions with conotruncal cardiac malformations [Goldmuntz et al., 1993], FISH analysis with the N25 probe (ONCOR) was performed. The patient was found to have a 22q11.2 microdeletion.

The patient's mother is in good health, however she did not speak clearly until age 3 and continues to have hypernasal speech. She also has a history of a repaired umbilical hernia, mitral valve prolapse, and bilateral placement of myringotomy tubes. Deletion studies demonstrated a maternal 22q11.2 deletion. The patient's maternal grandmother is in good health and does not have a 22q11.2 deletion by FISH. She had one spontaneous abortion at 8 weeks gestation and 4 term deliveries. The patient's maternal grandfather died at age 40 years of pancreatic cancer; he had previously been healthy. The patient's maternal uncle had a daughter with aortic stenosis. Deletion studies have not yet been performed in these family members.

PATIENT 2

This young man presented at age 18 years for correction of his broad, bifid nasal tip. At this time he was found to have a submucous cleft palate, bifid uvula and velopharyngeal incompetence. Neonatally he had hypocalcemia, atrial septal defect and delayed development. At 28 years he was evaluated by a neurologist for dysmetria and bradykinesia caused by cerebellar atrophy [Lynch et al., 1995]. His height was 185 cm (90th centile), OFC was 57.5 cm (75–90th centile). Eyes were deep set, inner canthal distance was 3 cm, outer canthal distance was 9.5 cm and inter pupillary distance was 5.8 cm (75–90th centile). He had a prominent nasal root with a bulbous, dimpled tip, and prominent ears with thick, overfolded helices (Fig. 2). His speech was dysarthric and his gait wide based. Because of the cleft

Contract grant sponsor: The National Institute on Deafness and Other Communicative Disorder, National Institute of Health; Contract grant number: DC 02027.

*Correspondence to: Karen W. Gripp, Clinical Genetics, Children's Hospital of Philadelphia, 34th and Civic Center Blvd., Philadelphia, PA, 19104.

Received 2 February 1996; Accepted 31 July 1996



Fig. 1. Patient 1 at age 5 weeks; note broad nose with bifid tip.

palate, developmental delay, neonatal hypocalcemia and the facial anomalies the diagnosis of velo-cardio-facial syndrome (VCFS) was suspected. Cytogenetic and FISH analysis demonstrated an interstitial deletion of chromosome 22q11.2.

DISCUSSION

The 22q11.2 microdeletion syndrome can present as DiGeorge syndrome [Driscoll et al., 1992a; Wilson et al., 1992], as VCFS [Driscoll et al., 1992b; Kelly et al., 1993], and as conotruncal anomaly face syndrome [Burn et al., 1993]. These syndromes share characteristic facial findings including broad nasal root and prominent nasal tip, hypertelorism, small mouth, and posteriorly angulated, prominent ears with a thickened helix. Congenital conotruncal cardiac defects, cleft palate, and thymic and parathyroid hypoplasia or aplasia

can also occur in these conditions. Due to the great variability of findings in affected patients a high index of suspicion is frequently required to make the diagnosis, as is demonstrated by the fact that some patients are identified only after a close relative was found to have the deletion. The presence of multiple, including subtle facial, anomalies can be helpful in diagnosing patients in a timely fashion.

We suggest that the occurrence of a nasal dimple in our two patients with 22q11.2 deletion is not coincidental. The dimpled appearance of the nasal tip probably represents the extreme of the broad and bulbous tip usually seen in this syndrome (see published photographs of case 4 and case 14 in Jedelev et al., 1992; mother of family 1 in De Silva et al., 1995). It should not be confused with the bifid nose in frontonasal dysplasia [Sedano et al., 1970].




Fig. 2. Patient 2 at age 10 years (left) and at age 18 years (right); note broad, bifid nasal tip and protruding ears.

De Moor et al. [1987] reported 3 patients with TOF, frontonasal dysplasia and medial nasal grooves, similar in appearance to those of our patients. Their growth was below the 3rd centile, a more significant growth failure than expected for their cardiac condition and not typical for frontonasal dysplasia. One of them had facial findings seen in the 22q11.2 deletion syndrome, including a very broad nasal root with hypoplastic alae nasi and, as evident in the published photograph, protruding ears. We wondered if this patient might have a 22q11.2 deletion, but unfortunately he was unavailable for further studies.

In summary, we suggest that the finding of a dimpled nasal tip, in combination with other anomalies, may be consistent with the diagnosis of 22q11.2 deletion syndrome.

ACKNOWLEDGMENTS

We thank Peter Beighton, MD for his help in trying to locate the South African patients. This work is supported in part by DC 02027 from the National Institute on Deafness and other Communication Disorders, National Institute of Health.

REFERENCES

- Burn J, Takao A, Wilson D, Cross I, Momma K, Wadey R, Scambler P, Goodship J (1993): Conotruncal anomaly face syndrome is associated with a deletion within chromosome 22q11. *J Med Genet* 30:822–824.
- De Moor MM, Baruch R, Human DG (1987): Frontonasal dysplasia associated with tetralogy of Fallot. *J Med Genet* 24:107–109.
- De Silva D, Duffy P, Booth P, Auchterlonie I, Morrison N, Dean JCS (1995): Family studies in chromosome 22q11 deletion: further demonstration of phenotypic heterogeneity. *Clin Dysmorphol* 4:294–303.
- Driscoll DA, Budarf ML, Emanuel BS (1992a): A genetic etiology for DiGeorge syndrome: consistent deletions and microdeletions of 22q11. *Am J Hum Genet* 50:924–933.
- Driscoll DA, Spinner NB, Budarf ML, McDonald-McGinn DM, Zackai EH, Goldberg RB, Shprintzen RJ, Saal HM, Zonana J, Jones MC, Emanuel BS (1992b): Deletions and microdeletions of 22q11.2 in velo-cardio-facial syndrome. *Am J Med Genet* 44:261–268.
- Goldmuntz E, Driscoll DA, Budarf ML, Zackai EH, McDonald-McGinn DM, Biegel JA, Emanuel BS (1993): Microdeletions of chromosomal region 22q11 in patients with congenital conotruncal cardiac defects. *J Med Genet* 30:807–812.
- Jedele KB, Michels VV, Puga FJ, Feldt RH (1992): Velo-cardio-facial syndrome associated with ventricular septal defect, pulmonary atresia, and hypoplastic pulmonary arteries. *Pediatrics* 89:915–919.
- Kelly D, Goldberg R, Wilson D, Lindsay E, Carey A, Goodship J, Burn J, Cross I, Shprintzen RJ, Scambler PJ (1993): Confirmation that the velo-cardio-facial syndrome is associated with haplo-insufficiency of genes at chromosome 22q11. *Am J Med Genet* 45:308–312.
- Lynch DR, McDonald-McGinn DM, Zackai EH, Emanuel BS, Driscoll DA, Whitaker LA, Fischbek KH (1995): Cerebellar atrophy in a patient with velocardiofacial syndrome. *J Med Genet* 32:561–563.
- Sedano HO, Cohen MMJ, Jirasek J, Gorlin RJ (1970): Frontonasal dysplasia. *J Pediatr* 76:906–913.
- Wilson DI, Cross IE, Goodship JA, Brown J, Scambler PJ, Bain HH, Taylor JF, Walsh K, Bankier A, Burn J, et al (1992): A prospective cytogenetic study of 36 cases of DiGeorge syndrome. *Am J Hum Genet* 51:957–963.